

Global Behavior of Solutions to Two Classes of Second Order Rational Difference Equations

Sukanya Basu Orlando Merino
 University of Rhode Island, Kingston, RI 02881
sukanya@math.uri.edu merino@math.uri.edu

Abstract

For nonnegative real numbers $\alpha, \beta, \gamma, A, B$ and C such that $B + C > 0$ and $\alpha + \beta + \gamma > 0$, the difference equation

$$x_{n+1} = \frac{\alpha + \beta x_n + \gamma x_{n-1}}{A + B x_n + C x_{n-1}}, \quad n = 0, 1, 2, \dots$$

has a unique positive equilibrium. A proof is given here for the following statements:

Theorem 1. *For every choice of positive parameters $\alpha, \beta, \gamma, A, B$ and C , all solutions to the difference equation*

$$x_{n+1} = \frac{\alpha + \beta x_n + \gamma x_{n-1}}{A + B x_n + C x_{n-1}}, \quad n = 0, 1, 2, \dots, \quad x_{-1}, x_0 \in [0, \infty)$$

converge to the positive equilibrium or to a prime period-two solution.

Theorem 2. *For every choice of positive parameters $\alpha, \beta, \gamma, A, B$ and C , all solutions to the difference equation*

$$x_{n+1} = \frac{\alpha + \beta x_n + \gamma x_{n-1}}{B x_n + C x_{n-1}}, \quad n = 0, 1, 2, \dots, \quad x_{-1}, x_0 \in (0, \infty)$$

converge to the positive equilibrium or to a prime period-two solution.

Contents

1	Introduction and Main Results	2
2	Results from the literature	4
3	Existence of an Invariant And Attracting Interval	5
4	The Equation (3-2) with $r \geq 0, p > q$ and $\frac{qr}{p-q} < \frac{p}{q}$	9
5	Proof of Theorem 3	15
6	Proof of Theorem 4	19
7	Appendix: Computer Algebra System Code	21

Key Words: difference equation, rational, global behavior, global attractivity, period-two solution.

AMS 2000 Mathematics Subject Classification: Primary: 39A05 Secondary: 39A11

1 Introduction and Main Results

In their book [11], M. Kulenović and G. Ladas initiated a systematic study of the difference equation

$$x_{n+1} = \frac{\alpha + \beta x_n + \gamma x_{n-1}}{A + B x_n + C x_{n-1}}, \quad n = 0, 1, 2, \dots \quad (1)$$

for nonnegative real numbers $\alpha, \beta, \gamma, A, B$ and C such that $B+C > 0$ and $\alpha+\beta+\gamma > 0$, and for nonnegative or positive initial conditions x_{-1}, x_0 . Under these conditions, (1) has a unique positive equilibrium. One of their main ideas in this undertaking was to make the task more manageable by considering separate cases when one or more of the parameters in (1) is zero. The need for this strategy is made apparent by cases such as the well known *Lyness Equation* [17], [22], [9].

$$x_{n+1} = \frac{\alpha + x_n}{x_{n-1}} \quad (2)$$

whose dynamics differ significantly from other equations in this class. There are a total of 42 cases that arise from (1) in the manner just discussed, under the hypotheses $B + C > 0$ and $\alpha + \beta + \gamma > 0$. The recent publications [1], [2] give a detailed account of the progress up to 2007 in the study of dynamics of the class of equations (1). After a sustained effort by many researchers (for extensive references, see [1], [2]), there are some cases that have resisted a complete analysis. We list them below in normalized form, as presented in [1], [2].

$$x_{n+1} = \frac{\alpha + x_n}{A + x_{n-1}} \quad (3)$$

$$x_{n+1} = \frac{\alpha + x_n}{x_n + C x_{n-1}} \quad (4)$$

$$x_{n+1} = \frac{\alpha + \beta x_n + \gamma x_{n-1}}{x_{n-1}} \quad (5)$$

$$x_{n+1} = \frac{\alpha + x_n}{A + B x_n + x_{n-1}} \quad (6)$$

$$x_{n+1} = \frac{\beta x_n + x_{n-1}}{A + B x_n + x_{n-1}} \quad (7)$$

$$x_{n+1} = \frac{\alpha + \beta x_n + \gamma x_{n-1}}{A + x_{n-1}} \quad (8)$$

$$x_{n+1} = \frac{\alpha + x_n + \gamma x_{n-1}}{B x_n + x_{n-1}} \quad (9)$$

$$x_{n+1} = \frac{\alpha + \beta x_n + x_{n-1}}{A + B x_n + x_{n-1}} \quad (10)$$

The dynamics of Equation (7) has been settled recently in [4], [20]. Global attractivity of the positive equilibrium of Equation (3) has been proved recently in [21]. Since Eq.(6) can be reduced to Eq.(3) through a change of variables [12], global behavior of solutions to (6) is also settled. Equation (5) is another equation that can be reduced to (3), through the change of variables $x_n = y_n + \gamma$ [10].

Ladas and co-workers [11], [1], [2], have posed a series of conjectures on these equations. One of them is the following.

Conjecture [Ladas et al.] *For equations (9) and (10), every solution converges to the positive equilibrium or to a prime period-two solution.*

In this article, we prove this conjecture. Our main results are the following.

Theorem 1 For every choice of positive parameters $\alpha, \beta, \gamma, A, B$ and C , all solutions to the difference equation

$$x_{n+1} = \frac{\alpha + \beta x_n + \gamma x_{n-1}}{A + B x_n + C x_{n-1}}, \quad n = 0, 1, 2, \dots, \quad x_{-1}, x_0 \in [0, \infty) \quad (3-3)$$

converge to the positive equilibrium or to a prime period-two solution.

Theorem 2 For every choice of positive parameters $\alpha, \beta, \gamma, A, B$ and C , all solutions to the difference equation

$$x_{n+1} = \frac{\alpha + \beta x_n + \gamma x_{n-1}}{B x_n + C x_{n-1}}, \quad n = 0, 1, 2, \dots, \quad x_{-1}, x_0 \in (0, \infty) \quad (11)$$

converge to the positive equilibrium or to a prime period-two solution.

A reduction of the number of parameters of Eq.(11) is obtained with the change of variables $x_n = \frac{\gamma}{C} y_n$, which yields the equation

$$y_{n+1} = \frac{r + p y_n + y_{n-1}}{q y_n + y_{n-1}}, \quad n = 0, 1, 2, \dots, \quad y_{-1}, y_0 \in (0, \infty) \quad (3-2)$$

where $r = \frac{\alpha C}{\gamma^2}$, $p = \frac{\beta}{\gamma}$, and $q = \frac{B}{C}$.

The number of parameters of Eq.(3-3) can also be reduced, which we proceed to do next. Consider the following affine change of variables which is helpful to reduce number of parameters and simplify calculations:

$$x_n = \left(\frac{\gamma}{C} + \frac{A}{B+C} \right) y_n - \frac{A}{B+C}, \quad (12)$$

With (12), Eqn.(3-3) may now be rewritten as

$$y_{n+1} = \frac{r + p y_n + y_{n-1}}{q y_n + y_{n-1}}, \quad n = 0, 1, 2, \dots, \quad y_{-1}, y_0 \in [L, \infty) \quad (3-2-L)$$

where

$$\begin{aligned} p &= \frac{AB + (B+C)\beta}{AC + (B+C)\gamma} \\ q &= \frac{B}{C}, \\ r &= \frac{C(B+C)(B\alpha + C\alpha - A\beta - A\gamma)}{(\gamma(B+C) + AC)^2}, \\ L &= \frac{AC}{AC + (B+C)\gamma} \end{aligned} \quad (13)$$

Theorems 1 and 2 can be reformulated in terms of the parameters p, q and r as follows.

Theorem 3 Let $\alpha, \beta, \gamma, A, B$ and C be positive numbers, and let p, q, r and L be given by relations (13). Then every solution to Eqn.(3-2-L) converges to the unique equilibrium or to a prime period-two solution.

Theorem 4 Let p, q, r be positive numbers. Then every solution to Eqn.(3-2) converges to the unique equilibrium or to a prime period-two solution.

In this paper we prove Theorems 3 and 4; Theorems 1 and 2 follow as an immediate corollary.

The two main differences between Eq.(3-2-L) and Eq.(3-2) are the set of initial conditions, and the possibility of having a negative value of r in Eq.(3-2-L), while only positive values of r are allowed in Eq.(3-2). Nevertheless, for both Eq.(3-2-L) and Eq.(3-2) the unique equilibrium has the formula:

$$\bar{y} = \frac{p+1 + \sqrt{(p+1)^2 + 4r(q+1)}}{2(q+1)}$$

Although it is not possible to prove Theorem 1 as a simple corollary to Theorem 2, the changes of variables leading to Theorems 3 and 4 will result in proofs to the former theorems that are greatly simplified.

Our main results Theorem 1 and Theorem 2 imply that when prime period-two solutions to Eq.(3-3) or Eq.(3-2) do not exist, then the unique equilibrium is a global attractor. We have not treated here certain questions about the global dynamics of Eq.(3-3) and Eq.(3-2), such as the character of the prime period-two solutions to either equation, or even for more general rational second order equations, when such solutions exist. This matter will be treated in an upcoming article of the authors [3].

This work is organized as follows. The main results are stated in Section 1. Results from the literature which are used here are given in Section 2 for convenience. In Section 3, it is shown that either every solution to Eq.(3-2-L) converges to the equilibrium, or there exists an invariant and attracting interval I with the property that the function $f(x, y)$ associated with the difference equation is coordinate-wise strictly-monotonic on $I \times I$. In Section 4, a global convergence result is obtained for Eq.(3-2) over a specific range of parameters and for initial conditions in an invariant compact interval. Theorem 3 is proved in Section 5, and the proof of Theorem 4 is given in Section 6. Section 7 includes computer algebra system code for performing certain calculations that involve polynomials with a large number of terms (over 365,000 in one case). These computer calculations are used to support certain statements in Section 4. Finally, we refer the reader to [11] for terminology and definitions that concern difference equations.

2 Results from the literature

The results in this subsection are from the literature, and they are given here for easy reference. The first result is a reformulation of Theorems (1.4.5) — (1.4.8) in [11].

Theorem 5 ([13], [11]) *Suppose a continuous function $f : [a, b]^2 \rightarrow [a, b]$ satisfies one of i.–iv.:*

i. $f(x, y)$ is nondecreasing in x, y , and

$$\forall (m, M) \in [a, b]^2, \quad (f(m, m) = m \ \& \ f(M, M) = M) \implies m = M$$

ii. $f(x, y)$ is nonincreasing in x, y , and

$$\forall (m, M) \in [a, b]^2, \quad (f(m, m) = M \ \& \ f(M, M) = m) \implies m = M$$

iii. $f(x, y)$ is nonincreasing in x and nondecreasing in y , and

$$\forall (m, M) \in [a, b]^2, \quad (f(m, M) = M \ \& \ f(M, m) = m) \implies m = M$$

iv. $f(x, y)$ is nondecreasing in x and nonincreasing in y , and

$$\forall (m, M) \in [a, b]^2, \quad (f(M, m) = M \ \& \ f(m, M) = m) \implies m = M$$

Then $y_{n+1} = f(y_n, y_{n-1})$ has a unique equilibrium in $[a, b]$, and every solution with initial values in $[a, b]$ converges to the equilibrium.

The following result is Theorem A.0.8 in [11].

Theorem 6 Suppose a continuous function $f : [a, b]^3 \rightarrow [a, b]$ is nonincreasing in all variables, and

$$\forall (m, M) \in [a, b]^3, \quad (f(m, m, m) = M \ \& \ f(M, M, M) = m) \implies m = M$$

Then $y_{n+1} = f(y_n, y_{n-1}, y_{n-2})$ has a unique equilibrium in $[a, b]$, and every solution with initial values in $[a, b]$ converges to the equilibrium.

Theorem 7 ([5]) Let I be a set of real numbers and let $F : I \times I \rightarrow I$ be a function $F(u, v)$ which decreases in u and increases in v . Then for every solution $\{x_n\}_{n=-1}^\infty$ of the equation

$$x_{n+1} = F(x_n, x_{n-1}), \quad n = 0, 1, \dots \quad (14)$$

the subsequences $\{x_{2n}\}$ and $\{x_{2n+1}\}$ of even and odd terms do exactly one of the following:

- (i) They are both monotonically increasing.
- (ii) They are both monotonically decreasing.
- (iii) Eventually, one of them is monotonically increasing and the other is monotonically decreasing.

Theorem 7 has this corollary.

Corollary 1 ([5]) If I is a compact interval, then every solution of Eq.(14) converges to an equilibrium or to a prime period-two solution.

Theorem 8 ([8]) Assume the following conditions hold:

- (i) $h \in C[(0, \infty) \times (0, \infty), (0, \infty)]$.
- (ii) $h(x, y)$ is decreasing in x and strictly decreasing in y .
- (iii) $x h(x, x)$ is strictly increasing in x .
- (iv) The equation

$$x_{n+1} = x_n h(x_n, x_{n-1}), \quad n = 0, 1, \dots \quad (15)$$

has a unique positive equilibrium $\bar{x}$.

Then $\bar{x}$ is a global attractor of all positive solutions of Eq.(15).

3 Existence of an Invariant And Attracting Interval

In this section we prove a proposition which is key for later developments. We will need the function

$$f(x, y) := \frac{r + px + y}{qx + y}, \quad x, y \in [L, \infty) \quad (16)$$

associated to Eq.(3-2-L).

Proposition 1 At least one of the following statements is true:

(A) Every solution to (3-2-L) converges to the equilibrium.

(B) There exist m^*, M^* with $L < m^* < M^*$ s.t.

- (i) $[m^*, M^*]$ is an invariant interval for Eq.(3-2-L), i.e., $f([m^*, M^*] \times [m^*, M^*]) \subset [m^*, M^*]$.
- (ii) Every solution to Eq.(3-2-L) eventually enters $[m^*, M^*]$.
- (iii) $f(x, y)$ is coordinate-wise strictly monotonic on $[m^*, M^*]^2$.

The next lemma states that the function $f(\cdot, \cdot)$ associated to Eq.(3-2-L) is bounded.

Lemma 1 *There exist positive constants $\mathcal{L}$ and $\mathcal{U}$ such that $L < \mathcal{L}$ and*

$$\mathcal{L} \leq f(x, y) \leq \mathcal{U}, \quad x, y \in [L, \infty) \quad (17)$$

In particular,

$$f([\mathcal{L}, \mathcal{U}] \times [\mathcal{L}, \mathcal{U}]) \subset [\mathcal{L}, \mathcal{U}] \quad (18)$$

Proof. The function

$$\tilde{f}(x, y) = \frac{\alpha + \beta x + \gamma y}{A + Bx + Cy}, \quad (x, y) \in (0, \infty)^2$$

associated to Eq.(3-3) is bounded:

$$\frac{\min\{\alpha, \beta, \gamma\}}{\max\{A, B, C\}} \leq \frac{\alpha + \beta x + \gamma y}{A + Bx + Cy} \leq \frac{\max\{\alpha, \beta, \gamma\}}{\min\{A, B, C\}}, \quad (x, y) \in (0, \infty)^2$$

Set $\tilde{\mathcal{L}} := \frac{\min\{\alpha, \beta, \gamma\}}{\max\{A, B, C\}}$ and $\tilde{\mathcal{U}} := \frac{\max\{\alpha, \beta, \gamma\}}{\min\{A, B, C\}}$. The affine change of coordinates (12) maps the rectangular region $[\tilde{\mathcal{L}}, \tilde{\mathcal{U}}]^2$ onto a rectangular region $[\mathcal{L}, \mathcal{U}]^2$ which satisfies (17) and (18). $\square$

Lemma 2 *If $p = q$, then every solution to Eq.(3-2-L) converges to the unique equilibrium.*

Proof. If $p = q$ then $D_1 f(x, y) = -\frac{pr}{(px+y)^2}$ and $D_2 f(x, y) = -\frac{r}{(px+y)^2}$. Thus, depending on the sign of r , the function $f(x, y)$ is either nondecreasing in both coordinates, or nonincreasing in both coordinates on $[L, \infty)$. By Lemma 1, all solutions $\{y_n\}_{n=-1}^\infty$ satisfy $y_n \in [\mathcal{L}, \mathcal{U}]$ for $n \geq 1$. A direct algebraic calculation may be used to show that all solutions $(m, M) \in [\mathcal{L}, \mathcal{U}]$ of either one of the systems of equations

$$\begin{cases} M &= f(M, M) \\ m &= f(m, m) \end{cases} \quad \text{and} \quad \begin{cases} M &= f(m, m) \\ m &= f(M, M) \end{cases}$$

necessarily satisfy $m = M$. In either case, the hypotheses (i) or (ii) of Theorem 5 are satisfied, and the conclusion of the lemma follows. $\square$

We will need the following elementary result, which is given here without proof.

Lemma 3 *Suppose $q \neq p$. The function $f(x, y)$ has continuous partial derivatives on $(L, \infty)^2$, and*

- i. $D_1 f(x, y) = 0$ if and only if $y = \frac{qr}{q-p}$, and $D_1 f(x, y) > 0$ if and only if $(p - q)y > qr$.
- ii. $D_2 f(x, y) = 0$ if and only if $x = \frac{-r}{p-q}$, and $D_2 f(x, y) > 0$ if and only if $(q - p)x > r$.

We will need to refer to the values K_1 and K_2 where the partial derivatives of $f(x, y)$ change sign.

Definition 1 If $p \neq q$, set

$$K_1 := \frac{qr}{p-q} \quad \text{and} \quad K_2 := \frac{-r}{p-q}$$

Definition 2 For $L \leq m \leq M$, let

$$\phi(m, M) := \min\{f(x, y) : (x, y) \in [m, M]^2\} \quad \text{and} \quad \Phi(m, M) := \max\{f(x, y) : (x, y) \in [m, M]^2\}$$

Lemma 4 Suppose $p \neq q$. If $[m, M] \subset [\mathcal{L}, \mathcal{U}]$ is an invariant interval for Eq.(3-2-L) with $m \leq K_1 \leq M$ or $m \leq K_2 \leq M$, then $m < \phi(m, M)$ or $\Phi(m, M) < M$ or $m = M = \bar{y}$.

Proof. By definition of ϕ and Φ , $m \leq \phi(m, M)$ and $\Phi(m, M) \leq M$. Suppose

$$m = \phi(m, M) \quad \text{and} \quad \Phi(m, M) = M \tag{19}$$

The proof will be complete when it is shown that $m = M$. There are a total of four cases to consider: (a) $r \geq 0$ and $p > q$, (b) $r < 0$ and $p < q$, (c) $r \geq 0$ and $p < q$, and (d) $r < 0$ and $p > q$. We present the proof of case (a) only, as the proof of the other cases is similar.

If $r \geq 0$ and $p > q$, then $K_1 \in [m, M]$ and $K_2 \notin [m, M]$. Note that

$$[m, M] \times [m, M] = [m, M] \times [m, K_1] \cup [m, M] \times [K_1, M].$$

By Lemma 3, the signs of the partial derivatives of $f(x, y)$ are constant on the interior of each of the sets $[m, M] \times [m, K_1]$ and $[m, M] \times [K_1, M]$, as shown in the diagram.




Figure 1: The arrows indicate type of coordinate-wise monotonicity of $f(x, y)$ on each region.

Since $f(x, y)$ is nonincreasing in both x and y on $[m, M] \times [m, K_1]$,

$$f(M, K_1) \leq f(x, y) \leq f(m, m) \quad \text{for } (x, y) \in [m, M] \times [m, K_1]. \tag{20}$$

Similarly, $f(x, y)$ is nondecreasing in x and nonincreasing in y on $[m, M] \times [K_1, M]$, hence

$$f(m, M) \leq f(x, y) \leq f(M, K_1) \quad \text{for } (x, y) \in [m, M] \times [K_1, M]. \tag{21}$$

From (20) and (21) one has

$$\phi(m, M) = f(m, M) \quad \text{and} \quad \Phi(m, M) = f(m, m) \tag{22}$$

Combine (22) with relation (19) to obtain the system of equations

$$\begin{cases} f(m, M) = m \\ f(m, m) = M \end{cases} \tag{23}$$

Eliminating M from system (23) gives the cubic in m

$$q(q+1)m^3 + (1-pq)m^2 + (-1-p-qr)m - r = 0 \quad (24)$$

which has the roots

$$-\frac{1}{q}, \quad \frac{1-p-\sqrt{(1+p)^2+4r(1+q)}}{2(1+q)}, \quad \text{and} \quad \frac{1-p+\sqrt{(1+p)^2+4r(1+q)}}{2(1+q)} \quad (25)$$

Only one root in the list (25) is positive, namely

$$m = \frac{1-p+\sqrt{(1+p)^2+4r(1+q)}}{2(1+q)} = \bar{y}$$

Substituting into one of the equations of system (23) one also obtains $M = \bar{y}$, which gives the desired relation $m = M = \bar{y}$. $\square$

Definition 3 Let $m_0 := \mathcal{L}$, $M_0 := \mathcal{U}$, and for $\ell = 0, 1, 2, \dots$ let $m_\ell := \phi(m_\ell, M_\ell)$, $M_\ell := \Phi(m_\ell, M_\ell)$.

By the definitions of m_ℓ , M_ℓ , $\phi(\cdot, \cdot)$ and $\Phi(\cdot, \cdot)$, we have that $[m_{\ell+1}, M_{\ell+1}] \subset [m_\ell, M_\ell]$ for $\ell = 0, 1, 2, \dots$. Thus the sequence $\{m_\ell\}$ is nondecreasing and $\{M_\ell\}$ is nonincreasing. Let $m^* := \lim m_\ell$ and $M^* := \lim M_\ell$.

Lemma 5 Suppose $p \neq q$. Either there exists $N \in \mathbb{N}$ such that $\{K_1, K_2\} \cap [m_N, M_N] = \emptyset$, or $m^* = M^* = \bar{y}$.

Proof. Arguing by contradiction, suppose $m^* < M^*$ and for all $\ell \in \mathbb{N}$, $\{K_1, K_2\} \cap [m_\ell, M_\ell] \neq \emptyset$. Since the intervals $[m_\ell, M_\ell]$ are nested and $\cap [m_\ell, M_\ell] = [m^*, M^*]$, it follows that $\{K_1, K_2\} \cap [m^*, M^*] \neq \emptyset$. By Lemma 4, we have

$$m^* < \phi(m^*, M^*) \quad \text{or} \quad \Phi(m^*, M^*) < M^* \quad (26)$$

Continuity of the functions ϕ and Φ implies

$$\phi(m^*, M^*) = \lim \phi(m_\ell, M_\ell) = \lim m_{\ell+1} = m^* \quad \text{or} \quad \Phi(m^*, M^*) = \lim \Phi(m_\ell, M_\ell) = \lim M_{\ell+1} = M^* \quad (27)$$

Statements (26) and (27) give a contradiction. $\square$

Proof of Proposition 1. Suppose statement (A) is not true. By Lemma 2, one must have $p \neq q$. Note that if $\{y_\ell\}$ is a solution to Eq.(3-2-L), then $y_{\ell+1} \in [m_\ell, M_\ell]$ for $\ell = 0, 1, 2, \dots$. If $m^* = M^*$, since $m_\ell \rightarrow m^*$ and $M_\ell \rightarrow M^*$ we have $y_\ell \rightarrow \bar{y}$, but this is statement (A) which we are negating. Thus $m^* < M^*$, and by Lemma 4 there exists $N \in \mathbb{N}$ such that $\{K_1, K_2\} \cap [m_N, M_N] = \emptyset$, so $f(x, y)$ is coordinate-wise monotonic on $[m_N, M_N]$. The set $[m_N, M_N]$ is invariant, and every solution enters $[m_N, M_N]$ starting at least with the term with subindex $N+1$. We have shown that if statement (A) is not true, then statement (B) is necessarily true. This completes the proof of the proposition. $\square$

4 The Equation (3-2) with $r \geq 0$, $p > q$ and $\frac{qr}{p-q} < \frac{p}{q}$

In this section we restrict our attention to the equation

$$y_{n+1} = f(x_n, x_{n-1}) \quad n = 0, 1, \dots, \quad x_{-1}, x_0 \in (0, \infty) \quad (3-2)$$

where

$$f(x, y) := \frac{r + px + y}{qx + y},$$

For $p > 0$, $q > 0$, and $r \geq 0$, Eq.(3-2) has a unique *positive* equilibrium

$$\bar{y} = \frac{p + 1 + \sqrt{(p + 1)^2 + 4r(q + 1)}}{2(q + 1)} \quad (28)$$

We note that if $I \subset (0, \infty)$ is an invariant compact interval, then necessarily $\bar{y} \in I$.

The goal in this section is to prove the following proposition, which will provide an important part of the proofs of Theorems 1 and 4.

Proposition 2 *Let p, q and r be real numbers such that*

$$p > q > 0, \quad r \geq 0, \quad \text{and} \quad \frac{qr}{p-q} < \frac{p}{q} \quad (29)$$

and let $[\tilde{m}, \tilde{M}] \subset (\frac{qr}{p-q}, \frac{p}{q})$ be a compact invariant interval for Eq.(3-2). Then every solution to Eq.(3-2) with $x_{-1}, x_0 \in [\tilde{m}, \tilde{M}]$ converges to the equilibrium.

Proposition 2 follows from Lemmas 6, 7 and 9, which are stated and proved next.

Lemma 6 *Assume the hypotheses to Proposition 2. If either $q \geq 1$ or $p \leq 1$, then every solution to Eq.(3-2) with $x_{-1}, x_0 \in [\tilde{m}, \tilde{M}]$ converges to the equilibrium.*

Proof. We verify that hypothesis (iv) of Theorem 5 is true. Since $x > \frac{rq}{p-q}$ for $x \in [\tilde{m}, \tilde{M}]$, the function $f(x, y)$ is increasing in x and decreasing in y for $(x, y) \in [\tilde{m}, \tilde{M}]^2$ by Lemma 3. Let $m, M \in [\tilde{m}, \tilde{M}]$ be such that $m \neq M$ and

$$\begin{cases} f(M, m) - M = 0 \\ f(m, M) - m = 0 \end{cases} \quad (30)$$

We show first that system (30) has no solutions if either $q \geq 1$ or $p \leq 1$. By eliminating denominators in both equations in (30),

$$\begin{cases} m - mM + Mp - M^2q + r = 0 \\ M - mM + mp - m^2q + r = 0 \end{cases} \quad (31)$$

and by subtracting terms in (31) one obtains

$$(M - m)(1 - p + q(m + M)) = 0 \quad (32)$$

Since $m \neq M$, we have $q(m + M) = p - 1$, which implies that for $p \leq 1$ there are no solutions to system (30) which have both coordinates positive. Now assume $p > 1$; from (32), $m = \frac{p-1-Mq}{q}$, and substitute the latter into (30) to see that $x = M$ is a solution to the quadratic equation

$$(1 - q)x^2 + \frac{(-1 + p)(-1 + q)}{q}x + \frac{-1 + p + qr}{q} = 0 \quad (33)$$

By a symmetry argument, one has that $x = m$ is also a solution to (33). By inspection of the coefficients of the polynomial in the left-hand-side of (33) one sees that two positive solutions are possible only when $q < 1$. To get the conclusion of the lemma, note that the fact that (30) has no solutions with $m \neq M$ is just hypothesis (iv) of Theorem 5. $\square$

Lemma 7 *Assume the hypotheses to Proposition 2. If*

$$r \leq p^2 q - p \quad (34)$$

then every solution to Eq.(3-2) with $x_{-1}, x_0 \in [\tilde{m}, \tilde{M}]$ converges to the equilibrium.

Proof. By substituting $x_n = f(x_{n-1}, x_{n-2})$ into $x_{n+1} = f(x_n, x_{n-1})$ we obtain

$$x_{n+1} = \frac{r + p x_n + x_{n-1}}{q x_n + x_{n-1}} = \frac{r + p \frac{r + p x_{n-1} + x_{n-2}}{q x_{n-1} + x_{n-2}} + x_{n-1}}{q \frac{r + p x_{n-1} + x_{n-2}}{q x_{n-1} + x_{n-2}} + x_{n-1}}, \quad n = 0, 1, \dots \quad (35)$$

that is,

$$x_{n+1} = \hat{f}(x_n, x_{n-1}, x_{n-2}), \quad \text{where } \hat{f}(x, y, z) = \frac{p r + p^2 y + q r y + q y^2 + p z + r z + y z}{q r + p q y + q x^2 + q z + y z} \quad (36)$$

where the x has been kept in $\hat{f}(x, y, z)$ for bookkeeping purposes. Thus $\hat{f}(x, y, z)$ is constant in x . We claim $\hat{f}(x, y, z)$ is decreasing in both y and z . To see that the partial derivative

$$D_3 \hat{f}(x, y, z) = - \frac{(r + (p - q) y) (-q r + (p - q) y)}{(q r + p q y + q y^2 + q z + y z)^2} \quad (37)$$

is negative just use $p > q$ and the inequality $(p - q) y - q r > 0$, which is true by Lemma 3. The remaining partial derivative is

$$D_2 \hat{f}(x, y, z) = - \frac{L(y, z)}{(q r + p q y + q y^2 + q z + y z)^2} \quad (38)$$

where

$$h(y, z) := -q^2 r^2 + 2 p q r y - 2 q^2 r y + p^2 q y^2 - p q^2 y^2 + q^2 r y^2 + p r z - q r z + p q r z - q^2 r z + 2 p q y z - 2 q^2 y z + 2 q r y z + p z^2 - q z^2 + r z^2$$

We have,

$$D_1 h(y, z) = 2 (p - q) q r + 2 q (p^2 - p q + q r) y + 2 q (p - q + r) z > 0, \quad (39)$$

$$D_2 h(y, z) = (p - q) (1 + q) r + 2 q (p - q + r) y + 2 (p - q + r) z > 0 \quad (40)$$

Since $\frac{q r}{p - q} \leq \tilde{m}$,

$$h(y, z) \geq h\left(\frac{q r}{p - q}, \frac{q r}{p - q}\right) = \frac{q (1 + q)^2 r^2 (p^2 - p q + q r)}{(p - q)^2} > 0, \quad y, z \in [\tilde{m}, \tilde{M}]$$

thus we conclude that $D_2 \hat{f}(x, y, z) < 0$ for $x, y, z \in [\tilde{m}, \tilde{M}]$.

To complete the proof we verify the hypotheses of Theorem 6. We claim that the system of equations

$$\begin{cases} \hat{f}(m, m, m) - M &= 0 \\ \hat{f}(M, M, M) - m &= 0 \end{cases} \quad (41)$$

has no solutions (m, M) with $m \neq M$ whenever hypothesis (34) holds.

By eliminating denominators in both equations in (41) one obtains

$$\begin{aligned} -m^2 + m^2 M - m p - m p^2 - m^2 q + m M q + m^2 M q + m M p q - m r - p r - m q r + M q r &= 0 \\ -M^2 + m M^2 - M p - M p^2 + m M q - M^2 q + m M^2 q + m M p q - M r - p r + m q r - M q r &= 0 \end{aligned} \quad (42)$$

and by subtracting terms in (42) one obtains

$$((m - M) (-m - M + m M - p - p^2 - m q - M q + m M q - r - 2 q r)) = 0 \quad (43)$$

Since $m \neq M$, we may use the second factor in the left-hand-side term of (43) to solve for M in terms of m , which upon substitution into $\hat{f}(m, m, m) = M$ and simplification yields the equation

$$\frac{a_2 m^2 + a_1 m + a_0}{(-1 + m) (1 + q) (m^2 + m q + m^2 q + m p q + q r)} = 0 \quad (44)$$

where

$$\begin{aligned} a_0 &= r (p + 2 p q + p^2 q + q r + 2 q^2 r) \\ a_1 &= p + p^2 + 2 p q + 3 p^2 q + p^3 q + r - p r + 4 q r + 4 q^2 r + 2 p q^2 r \\ a_2 &= (1 + q) (1 + 2 q + p q + q r) \end{aligned}$$

By hypothesis (34) we have $r p \leq p^3 q - p^2 < p^3 q$, hence $p^3 q - r p > 0$, which implies $a_1 \geq 0$. By direct inspection one can see that $a_0 > 0$ and $a_2 > 0$. Thus (44) has no positive solutions, and we conclude that (41) has no solutions $(m, M) \in [\tilde{m}, \tilde{M}]$ with $m \neq M$. We have verified the hypotheses of Theorem 6, and the conclusion of the lemma follows. $\square$

Lemma 8 *Let $p > 0$, $q > 0$ and $r \geq 0$. If the positive equilibrium $\bar{y}$ of Eq.(3-2) satisfies $\bar{y} < \frac{p}{q}$, then $\bar{y}$ is locally asymptotically stable (L.A.S.).*

Proof. Solving for r in

$$\bar{y} = \frac{r + (p + 1) \bar{y}}{(q + 1) \bar{y}}$$

gives

$$r = (q + 1) \bar{y}^2 - (p + 1) \bar{y} \quad (45)$$

Then a calculation shows

$$D_1 f(\bar{y}, \bar{y}) = \frac{p - q \bar{y}}{\bar{y} (q + 1)}$$

$$D_2 f(\bar{y}, \bar{y}) = -\frac{\bar{y} - 1}{\bar{y} (q + 1)}$$

Set $t_1 := D_1 f(\bar{y}, \bar{y})$ and $t_2 := D_2 f(\bar{y}, \bar{y})$. The equilibrium $\bar{y}$ is locally asymptotically stable if the roots of the characteristic polynomial

$$\rho(x) = x^2 - t_1 x - t_2$$

have modulus less than one [11]. By the Schur-Cohn Theorem, $\bar{y}$ is L.A.S. if and only if $|t_1| < 1 - t_2 < 2$. It can be easily verified that $1 - t_2 < 2$ if and only if $0 < q \bar{y} + 1$ which is true regardless of the allowable parameter values. Since $p - q \bar{y} > 0$ by the hypothesis, we have $|t_1| = \left| \frac{p - q \bar{y}}{\bar{y} (q + 1)} \right| = \frac{p - q \bar{y}}{\bar{y} (q + 1)}$, hence some algebra gives $|t_1| < 1 - t_2$ if and only if

$$\frac{1}{2} \frac{p + 1}{q + 1} < \bar{y} \quad (46)$$

But (46) is a true statement by formula (28). We conclude $\bar{y}$ is L.A.S. $\square$

Lemma 9 Assume the hypotheses to Proposition 2. If

$$p > 1, \quad q < 1, \quad \text{and} \quad r > p^2 q - p \quad (47)$$

then every solution to Eq.(3-2) with $x_{-1}, x_0 \in [\tilde{m}, \tilde{M}]$ converges to the equilibrium.

Proof. The proof begins with a change of variable in Eq.(3-2) to produce a transformed equation with normalized coefficients analogous to those in the standard *normalized Lyness' Equation* [17], [22], [9]

$$z_{n+1} = \frac{\tilde{\alpha} + x_n}{x_{n-1}} \quad (48)$$

We seek to use an argument of proof similar to the one used in [21], in which one takes advantage of the existence of *invariant curves* of Lyness' Equation to produce a Lyapunov-like function for Eq.(3-2).

Set $y_n = pz_n$ in Eq.(3-2) to obtain the equation

$$z_{n+1} = \frac{a + z_n + g z_{n-1}}{b z_n + z_{n-1}}, \quad n = 0, 1, \dots \quad z_{-1}, z_0 \in (0, \infty) \quad (49)$$

where

$$a = \frac{r}{p^2}, \quad g = \frac{1}{p}, \quad b = q \quad (50)$$

We shall denote with $\bar{z}$ the unique equilibrium of Eq.(49). Note that

$$\bar{z} = p \bar{y} \quad (51)$$

It is convenient to parametrize Eq.(49) in terms of the equilibrium. We will use the symbol u to represent the equilibrium $\bar{z}$ of Eq.(49). By direct substitution of the equilibrium $u = \bar{z}$ into Eq.(49) we obtain

$$r = (b + 1) u^2 - (g + 1) u \quad (52)$$

By (52), $r \geq 0$ iff $u \geq \frac{g+1}{b+1}$. Using (52) to eliminate r from Eq.(49) gives the following equation for $b > 0$, $g > 0$ and $u \geq \frac{g+1}{b+1}$, equivalent to Eq.(49):

$$z_{n+1} = \frac{(b + 1) u^2 - (g + 1) u + z_n + g z_{n-1}}{b z_n + z_{n-1}}, \quad n = 0, 1, 2, \dots, \quad y_{-1}, y_0 \in (0, \infty) \quad (53)$$

Therefore it suffices to prove that all solutions of Eq.(53) converge to the equilibrium u .

The following statement is crucial for the proof of the proposition.

Claim 1 $u > 1$ if and only if $r > p^2 q - p$.

Proof. Since $\bar{y} = p \bar{z} = pu$, we have $u > 1$ if and only if $\bar{y} > p$, which holds if and only if

$$\frac{p + 1 + \sqrt{(p + 1)^2 + 4r(q + 1)}}{2(q + 1)} > p$$

After an elementary simplification, the latter inequality can be rewritten as $r > p^2 q - p$. □

By the hypotheses of the lemma, by Claim 1, and by (50) and (52) we have

$$b < 1, \quad g < 1, \quad 1 < u < \frac{1}{b}, \quad \text{and} \quad \frac{g + 1}{b + 1} \leq u \quad (54)$$

We now introduce a function which is the invariant function for (48) with constant $\tilde{\alpha} = u^2 - u$ (in this case the equilibrium of (48) is u):

$$g(x, y) = \frac{(1+x)(1+y)(u^2 - u + x + y)}{xy} \quad (55)$$

Note that $g(x, y) > 0$ for all $x, y \in (0, \infty)$ whenever $u > 1$. By using elementary calculus, one can show that the function $g(x, y)$ has a strict global minimum at (u, u) [9], [22], i.e.,

$$g(u, u) < g(x, y), \quad (x, y) \in (0, \infty)^2 \quad (56)$$

We need some elementary properties of the sublevel sets

$$S(c) := \{(s, t) \in (0, \infty) : g(s, t) \leq c\}, \quad c > 0$$

We denote with $Q_\ell(u, u)$, $\ell = 1, 2, 3, 4$ the four regions

$$\begin{aligned} Q_1(u, u) &:= \{(x, y) \in (0, \infty) \times (0, \infty) : u \leq x, u \leq y\} \\ Q_2(u, u) &:= \{(x, y) \in (0, \infty) \times (0, \infty) : x \leq u, u \leq y\} \\ Q_3(u, u) &:= \{(x, y) \in (0, \infty) \times (0, \infty) : x \leq u, y \leq u\} \\ Q_4(u, u) &:= \{(x, y) \in (0, \infty) \times (0, \infty) : u \leq x, y \leq u\} \end{aligned}$$

Let

$$T(x, y) := \left(y, \frac{(g+1)u^2 - (b+1)u + y + gx}{by + x} \right), \quad (x, y) \in (0, \infty) \times (0, \infty) \quad (57)$$

be the map associated to Eq. (53) (see [16]).

Claim 2 *If $(x, y) \in Q_2(u, u) \cup Q_4(u, u) \setminus \{(u, u)\}$, then $g(T(x, y)) < g(x, y)$.*

Proof. Set

$$\Delta_1(x, y) := g(x, y) - g(T(x, y)) \quad (58)$$

A calculation yields

$$\Delta_1(x, y) = -\frac{(1+x) F_1(x, y) F_2(x, y)}{xy (bx + y) F_3(x, y)} \quad (59)$$

where

$$\begin{aligned} F_1(x, y) &:= b(x - u)(y - \frac{1}{b}) + (y - u)(bu + y + u - g) \\ F_2(x, y) &:= b(x - u)^2 + b(x - u)u + b(x - u)u^2 + (u - y)(bu^2 + yg) \\ F_3(x, y) &:= (b+1)u^2 - (1+g)u + x + gy \end{aligned}$$

By (54), for $(x, y) \in Q_4(u, u) \setminus \{(u, u)\}$ we have $u \leq x$ and $y \leq u < \frac{1}{b}$ with $(x, y) \neq (u, u)$, therefore $F_1(x, y) < 0$, $F_2(x, y) > 0$ and $F_3(x, y) > 0$. Consequently $\Delta_1(x, y) > 0$ for $(x, y) \in Q_4(u, u) \setminus \{(u, u)\}$. To see that $\Delta_1(x, y) > 0$ for $(x, y) \in Q_2(u, u) \setminus \{(u, u)\}$ as well, rewrite $F_1(x, y)$ and $F_2(x, y)$ as follows:

$$\begin{aligned} F_1(x, y) &= b(u - x)(\frac{1}{b} - u) + (y - u)^2 + (y - u)u + (y - u)(u - g) + b(y - u)x \\ F_2(x, y) &= b(x - u)(x + u^2) - b(y - u)u^2 - (y - u)^2g - (y - u)ug \end{aligned}$$

For $(x, y) \in Q_2(u, u) \setminus \{(u, u)\}$ we have $x \leq u \leq y$ and $(x, y) \neq (u, u)$. Thus $F_1(x, y) > 0$, $F_2(x, y) < 0$, and $F_3(x, y) > 0$, which imply $\Delta_1(x, y) > 0$. $\square$

Claim 3 Suppose $g > b$. If $(x, y) \in Q_1(u, u) \cup Q_3(u, u) \setminus \{(u, u)\}$, then $g(T^2(x, y)) < g(x, y)$.

Proof. This proof requires extensive use of a computer algebra system to verify certain inequalities involving rational expressions. Here we give an outline of the steps, and refer the reader to Section 7 for the details.

Since $b < g < 1 < u < \frac{1}{b}$, and $\frac{g+1}{b+1} < u$ we may write

$$\begin{aligned} u &= \frac{g+1}{b+1} + t, \quad t > 0 \\ g &= \frac{b+\frac{s}{b}}{1+s}, \quad s > 0 \end{aligned} \tag{60}$$

The expression $\Delta_2 := g(x, y) - g(T^2(x, y))$ may be written as a single ratio of polynomials, $\Delta_2 = \frac{N}{D}$ with $D > 0$. The next step is to show $N > 0$ for $(x, y) \neq (u, u)$.

Points (x, y) in $Q_1(u, u)$ may be written in the form $x = u + v$, $y = u + w$, where $v, w \in [0, \infty)$. Substituting x, y, u and g in terms of v, w, s and t into the expression for N one obtains a rational expression $\frac{\tilde{N}}{D}$ with positive denominator. The numerator $\tilde{N}$ has some negative coefficients. At this points two cases are considered, $w \geq v$, and $w \leq v$. These can be written as $w = v + k$ and $v = w + k$ for nonnegative k . Substitution of each one of the latter expressions in $\tilde{N}$ gives a polynomial with positive coefficients. This proves $\Delta_2(x, y) > 0$ for $(x, y) \in Q_2(u, u)$.

If now we assume $(x, y) \in Q_3(u, u)$ with $(x, y) \neq (u, u)$, we may write

$$\begin{aligned} x &= \frac{u}{v+1} \quad v \in [0, \infty) \\ y &= \frac{u}{w+1} \quad w \in [0, \infty) \end{aligned} \tag{61}$$

The rest of the proof is as in the first case already discussed. Details can be found in Section 7. $\square$

Claim 4 Suppose $g < b$ and $u > 1$. If $(x, y) \in Q_1(u, u) \cup Q_3(u, u) \setminus \{(u, u)\}$, then $g(T^3(x, y)) < g(x, y)$.

Proof. The proof is analogous to the proof of Claim 3. We provide an outline. More details can be found in Section 7.

Since $1 < u$, we may write $u = 1 + t$ with $t > 0$. Also, $u < \frac{1}{b}$ implies $b < \frac{1}{u}$, and $b = \frac{1}{1+t+s}$ for $s > 0$. Since $g < b$ we may write $g = \frac{1}{1+t+s+\ell}$ for $\ell > 0$.

The expression $\Delta_3 := g(x, y) - g(T^3(x, y))$ may be written as a single ratio of polynomials, $\Delta_3 = \frac{N}{D}$ with $D > 0$. The next step is to show $N > 0$ for $(x, y) \neq (u, u)$. This is done in a way similar to the procedure described in Claim 3. $\square$

To complete the proof of the lemma, let $(\phi, \psi) \in (0, \infty) \times (0, \infty)$. Let $\{y_n\}_{n \geq -1}$ be the solution to (53) with initial condition $(y_{-1}, y_0) = (\phi, \psi)$, and let $\{T^n(\phi, \psi)\}_{n \geq 0}$ be the corresponding orbit of T . The following argument is essentially the same as the one found in [21]; we provided here for convenience. Define

$$\hat{c} := \liminf_n g(T^n(\phi, \psi)) \tag{62}$$

Note that $\hat{c} < \infty$, which can be shown by applying Claims 2, 3 and 4 repeatedly as needed to obtain a nonincreasing subsequence of $\{g(T^n(\phi, \psi))\}_{n \geq 0}$ that is bounded below by $g(u, u)$. Let $\{g(T^{n_k}(\phi, \psi))\}_{k \geq 0}$ be a subsequence convergent to $\hat{c}$. Therefore there exists $c > 0$ such that

$$g(T^{n_k}(\phi, \psi)) \leq c \quad \text{for all } k \geq 0,$$

that is,

$$T^{n_k}(\phi, \psi) \in S(c) := \{(s, t) : g(s, t) \leq c\} \quad \text{for } k \geq 0$$

The set $S(c)$ is closed by continuity of $g(x, y)$. Boundedness of $S(c)$ follows from

$$0 < x, y < \frac{(1+x)(1+y)(u^2 - u + x + y)}{xy} = g(x, y) = c, \quad \text{for } (x, y) \in S(c)$$

Thus $S(c)$ is compact, and there exists a convergent subsequence $\{T^{n_{k_\ell}}(\phi, \psi)\}_\ell$ with limit $(\hat{x}, \hat{y})$. Note that

$$\hat{c} = \lim_{\ell \rightarrow \infty} g(T^{n_{k_\ell}}(\phi, \psi)) = g(\hat{x}, \hat{y}) \quad (63)$$

We claim that $(\hat{x}, \hat{y}) = (u, u)$. If not, then by claims 2 3 and 4,

$$\min\{g(T(\hat{x}, \hat{y})), g(T^2(\hat{x}, \hat{y})), g(T^3(\hat{x}, \hat{y}))\} < \hat{c} \quad (64)$$

Let $\|\cdot\|$ denotes the euclidean norm. By (64) and continuity, there exists $\delta > 0$ such that

$$\|(s, t) - (\hat{x}, \hat{y})\| < \delta \implies \min\{g(T(s, t)), g(T^2(s, t)), g(T^3(s, t))\} < \hat{c} \quad (65)$$

Choose $L \in \mathbb{N}$ large enough so that

$$\|T^{n_{k_L}}(\phi, \psi) - (\hat{x}, \hat{y})\| < \delta \quad (66)$$

But then (65) and (66) imply

$$\min\{g(T^{n_{k_L}+1}(\phi, \psi)), g(T^{n_{k_L}+2}(s, t)), g(T^{n_{k_L}+3}(s, t))\} < \hat{c} \quad (67)$$

which contradicts the definition (62) of $\hat{c}$. We conclude $(\hat{x}, \hat{y}) = (u, u)$. From this and the definition of convergence of sequences we have that for every $\epsilon > 0$ there exists $L \in \mathbb{N}$ such that $\|T^{n_{k_L}}(\phi, \psi) - (u, u)\| < \epsilon$. Finally, since

$$\max\{|y_{n_{k_L}-1} - u|, |y_{n_{k_L}} - u|\} \leq \|(y_{n_{k_L}-1} - u, y_{n_{k_L}} - u)\| = \|T^{n_{k_L}}(\phi, \psi) - (u, u)\|$$

we have that for every $\epsilon > 0$ there exists $L \in \mathbb{N}$ such that $|y_{n_{k_L}} - u| < \epsilon$ and $|y_{n_{k_L}-1} - u| < \epsilon$. Since u is a locally asymptotically stable equilibrium for Eq.(53) by Lemma 8, it follows that $y_n \rightarrow u$. This completes the proof of the lemma. $\square$

5 Proof of Theorem 3

To prove Theorem 3 it is enough to assume statement (B) of Proposition 1. Also by Lemma 2 we may assume $p \neq q$ without loss of generality. Thus we make the following standing assumption, valid throughout the rest of this section for Eq.(3-2-L).

Standing Assumption (SA) Assume $p \neq q$ and that there exist m^*, M^* with $\mathcal{L} \leq m^* < M^* \leq \mathcal{U}$ such that for Eq.(3-2-L) and its associated function $f(x, y)$,

- (i) $[m^*, M^*]$ is an invariant interval.
- (ii) every solution eventually enters $[m^*, M^*]$.
- (iii) $f(x, y)$ is coordinate-wise strictly monotonic on $[m^*, M^*]^2$.

The function $f(x, y)$ is assumed to be coordinate-wise monotonic on $[m^*, M^*]$, and there are four possible cases in which this can happen: (a) $f(x, y)$ is increasing in both variables, (b) $f(x, y)$ is decreasing in both variables, (c) $f(x, y)$ is decreasing in x and increasing in y , and (d) $f(x, y)$ is increasing in x and decreasing in y .

We present several lemmas before completing the proof of Theorem 3.

By considering the restriction of the map T of Eq.(3-2-L) to $[m^*, M^*]^2$, an application of the Schauder Fixed Point Theorem [7] gives that $[m^*, M^*]^2$ contains the fixed point of T , namely $(\bar{y}, \bar{y})$. Thus we have the following result.

Lemma 10 $\bar{y} \in [m^*, M^*]$.

Lemma 11 *Neither one of the systems of equations*

$$(S_1) \quad \begin{cases} M &= f(M, M) \\ m &= f(m, m) \end{cases} \quad \text{and} \quad (S_2) \quad \begin{cases} M &= f(m, m) \\ m &= f(M, M) \end{cases}$$

have solutions $(m, M) \in [m^, M^*]^2$ with $m < M$.*

Proof. Since $x = \bar{y}$ is the only solution to $f(x, x) = x$, it is clear that only $(\bar{y}, \bar{y})$ satisfies (S_1) . Now let (m, M) be a solution to (S_2) . From straightforward algebra applied to $M - m = f(m, m) - f(M, M)$ one arrives at $(p + 1)(M - m) = 0$, which implies $m = M$. $\square$

Lemma 12 *Suppose $f(x, y)$ is increasing in x and decreasing in y for $(x, y) \in [m^*, M^*]$. Then $p - q > 0$.*

Proof. By the standing assumption (SA), $p \neq q$. By Lemma 3, the coordinate-wise monotonicity hypothesis, and the fact $\bar{y} \in [m^*, M^*]$ from Lemma 10, we have

$$(p - q)\bar{y} > q r \quad \text{and} \quad (q - p)\bar{y} < r \tag{68}$$

The inequalities in (68) cannot hold simultaneously unless $p - q > 0$. $\square$

Lemma 13 *If $f(x, y)$ is increasing in x and decreasing in y for $(x, y) \in [m^*, M^*]$, then $f([m^*, M^*]^2) \subset (1, \frac{p}{q})$.*

Proof. For $(x, y) \in [m^*, M^*]$, the function f is well defined and is componentwise strictly monotonic on the set $[x, \infty) \times [y, \infty)$. Then,

$$\begin{aligned} f(x, y) &< \lim_{s \rightarrow \infty} f(s, y) = \lim_{s \rightarrow \infty} \frac{r + p s + y}{q s + y} = \frac{p}{q} \\ f(x, y) &> \lim_{t \rightarrow \infty} f(x, t) = \lim_{t \rightarrow \infty} \frac{r + p x + t}{q x + t} = 1 \end{aligned} \tag{69}$$

$\square$

Lemma 14 *Let $p > 0$, $q > 0$ and $r \geq 0$. If $f(x, y)$ is increasing in x and decreasing in y on $[m^*, M^*]$, then*

$$\frac{q r}{p - q} < \frac{p}{q} \tag{70}$$

Proof. Since $\bar{y} \in [m^*, M^*]$ by Lemma 10, we have $D_1(\bar{y}, \bar{y}) > 0$ and $D_2(\bar{y}, \bar{y}) < 0$. By Lemma 12, $p > q$, and by Lemma 3,

$$(p - q) \bar{y} > q r \quad \text{and} \quad (q - p) \bar{y} < r \quad (71)$$

Then,

$$\bar{y} > \frac{q r}{p - q} \quad (72)$$

In addition, by Lemma 3,

$$\bar{y} = f(\bar{y}, \bar{y}) < \lim_{s \rightarrow \infty} f(s, \bar{y}) = \lim_{s \rightarrow \infty} \frac{r + p s + \bar{y}}{q s + \bar{y}} = \frac{p}{q} \quad (73)$$

□

Lemma 15 Suppose $f(x, y)$ is increasing in x and decreasing in y for $(x, y) \in [m^*, M^*]$. If $r < 0$, then $p - q + r > 0$.

Proof. Since $D_1 f(x, y) > 0$ for $(x, y) \in [m^*, M^*]$, and by Lemma 3, Lemma 10 and by Lemma 12, we have $\bar{y} > \frac{-r}{p-q}$, that is,

$$\sqrt{(p+1)^2 + 4r(q+1)} > \frac{-2r(q+1)}{p-q} - p - 1 \quad (74)$$

If the right-hand-side of inequality (74) is nonnegative, then, after squaring both sides of (74) we have

$$(p+1)^2 + 4r(q+1) > \left(\frac{-2r(q+1)}{p-q} \right)^2 + \frac{4r(q+1)(p+1)}{p-q} + (p+1)^2 \quad (75)$$

Further simplification of (75) and the hypothesis $r < 0$ yield

$$1 < \frac{r(q+1)}{(p-q)^2} + \frac{p+1}{p-q} \quad (76)$$

which, after some elementary algebra, implies $p - q + r > 0$. Now assume the right-hand-side of inequality (74) is negative, relation that we may rewrite as

$$\frac{-r}{p-q} < \frac{1}{2} \frac{p+1}{q+1} \quad (77)$$

If $\frac{1}{2} \frac{p+1}{q+1} \leq 1$, then $\frac{-r}{p-q} < 1$, which gives the conclusion $p - q + r > 0$. If $\frac{1}{2} \frac{p+1}{q+1} > 1$, that is, $p > 2q + 1$, then

$$p - q + r > q + r + 1 \quad (78)$$

Therefore if $q + r + 1 \geq 0$ the conclusion of the lemma follows from this and from (78). Assume

$$q + r + 1 < 0 \quad (79)$$

From relations (13) we have

$$q + r + 1 = \frac{(b+c)(a^2 c^2 + b c^2 \alpha + c^3 \alpha - a c^2 \beta + 2 a b c \gamma + a c^2 \gamma + b^2 \gamma^2 + 2 b c \gamma^2 + c^2 \gamma^2)}{c(a c + b \gamma + c \gamma)^2} \quad (80)$$

hence assumption (79) and relation (80) imply

$$R := a^2 c^2 + b c^2 \alpha + c^3 \alpha - a c^2 \beta + 2 a b c \gamma + a c^2 \gamma + b^2 \gamma^2 + 2 b c \gamma^2 + c^2 \gamma^2 < 0 \quad (81)$$

Further algebra gives

$$\frac{\gamma}{a c} R - (-c^2 \alpha + a c \gamma - c \beta \gamma + b \gamma^2) = c^2 \alpha + \frac{b c \alpha \gamma}{a} + \frac{c^2 \alpha \gamma}{a} + b \gamma^2 + c \gamma^2 + \frac{2 b \gamma^3}{a} + \frac{b^2 \gamma^3}{a c} + \frac{c \gamma^3}{a} > 0 \quad (82)$$

Since $R < 0$ by (81), from inequality (82) we have

$$-c^2 \alpha + a c \gamma - c \beta \gamma + b \gamma^2 < 0 \quad (83)$$

Finally, from (13) we have

$$p - q + r = -\frac{(b + c)^2 (-c^2 \alpha + a c \gamma - c \beta \gamma + b \gamma^2)}{c(a c + b \gamma + c \gamma^2)} \quad (84)$$

Combining (83) with (84) we obtain $p - q + r > 0$. $\square$

Lemma 16 *If $r < 0$ and $f(x, y)$ is increasing in x and decreasing in y for $(x, y) \in [m^*, M^*]$, then every solution converges to the equilibrium.*

Proof. Since $p - q + r > 0$ by Lemma 15, we have $K_2 = -\frac{r}{p-q} < 1$, which together with Lemma 13 implies that $[1, \frac{p}{q}]$ is an invariant, attracting compact interval such that $f(x, y)$ is increasing in x and decreasing in y on $[1, \frac{p}{q}]^2$. Since $f([1, \frac{p}{q}]^2) \subset (1, \frac{p}{q})$, we see that every solution to Eq.(3-2-L) eventually enters the invariant interval $(1, \frac{p}{q})$. The change of variables

$$y_n = \frac{1 + \frac{p}{q} z_n}{1 + z_n}, \quad \text{or} \quad z_n = \frac{x_n - 1}{\frac{p}{q} - x_n}, \quad (85)$$

transforms the equation

$$y_{n+1} = \frac{r + p y_n + y_{n+1}}{q y_n + y_{n+1}}, \quad n = 0, 1, \dots, \quad y_{-1}, y_0 \in \left(1, \frac{p}{q}\right) \quad (86)$$

into the equivalent equation

$$z_{n+1} = g(z_n, z_{n-1}), \quad n = 0, 1, \dots, \quad z_{-1}, z_0 \in (0, \infty) \quad (87)$$

where

$$g(w, v) := \frac{q(1+v)(-(q(p-q+r)) + (-p^2 + pq - qr)w)}{(1+w)(-q(p-q-qr) + (-p^2 + pq + q^2r)v)}$$

We claim that for $w, v \in (0, \infty)$, (a) $g(w, w)$ is increasing in w , (b) $g(w, v)/w$ is decreasing in w , and (c) $g(w, v)/w$ is decreasing in v . Indeed, since $p > q$, $r < 0$, $p - q + r > 0$, and $\frac{-r}{p-q} < \frac{p}{q}$ we have

$$\begin{aligned} \frac{d}{dw} (g(w, w)) &= \frac{-r q^2 (1+q) (-p+q)^2}{(-pq + q^2 + q^2 r - p^2 w + pq w + q^2 r w)^2} > 0 \\ \frac{\partial}{\partial w} \left(\frac{g(w, v)}{v} \right) &= -\frac{q(-p+q)^2 (q(p-q+r) + (p^2 - pq + qr)w)}{(-pq + q^2 + q^2 r - p^2 v + pq v + q^2 r v)^2 w (1+w)} < 0 \\ \frac{\partial}{\partial w} \left(\frac{g(w, v)}{w} \right) &= -\frac{q(1+v) (q(p-q+r) + 2q(p-q+r)w + (p^2 - pq + qr)w^2)}{(q(p-q-qr) + (p^2 - pq - q^2 r)v) w^2 (1+w)^2} < 0 \end{aligned}$$

Also, note that Eq.(87) has a unique equilibrium $\bar{z}$. Therefore hypotheses (1)–(4) of Theorem 8 are satisfied, so every solution $\{z_n\}$ to Eq.(87) converges to $\bar{z}$. By reversing the change of variables, one can conclude that every solution to Eq.(86) converges to the equilibrium. $\square$

Proof of Theorem 3. The four parts of the proof are:

- a. $f(x, y)$ is increasing in both x and y on $[m^*, M^*]^2$: By Lemma 11 the hypotheses of Theorem 5 part (i). is satisfied, hence every solution converges to the equilibrium $\bar{y}$.
- b. $f(x, y)$ is decreasing in both x and y on $[m^*, M^*]^2$: By Lemma 11 the hypotheses of Theorem 5 part (ii). is satisfied, hence every solution converges to the equilibrium $\bar{y}$.
- c. $f(x, y)$ is decreasing in x and increasing in y on $[m^*, M^*]^2$: By the corollary to Theorem 7 we conclude every solution converges to the unique equilibrium or to a prime period-two solution.
- d. $f(x, y)$ is increasing in x and decreasing in y on $[m^*, M^*]^2$: By Lemmas 3, 12, and 13, there is no loss of generality in assuming $[m^*, M^*] \subset (K, \frac{p}{q})$, where $K := \max\{\frac{-r}{p-q}, \frac{qr}{p-q}\}$, which we do. We consider two subcases. If $r \geq 0$, then Lemma 12, Lemma 14 and Proposition 2 imply that every solution converges to the unique equilibrium. If $r < 0$, then Lemma 16 implies that every solution converges to the unique equilibrium.

This completes the proof of Theorem 3. Since Theorem 3 is just a version of Theorem 1 obtained by an affine change of coordinates, we have also proved Theorem 1 as well. $\square$

6 Proof of Theorem 4

The first lemma guarantees solutions to Eq.(3-2) to be bounded.

Lemma 17 *Let $p > 0$, $q > 0$ and $r \geq 0$. There exist positive constants $\mathcal{L}$ and $\mathcal{U}$ such that every solution $\{x_n\}_{n=-1}^\infty$ to Eq.(3-2) satisfies $x_n \in [\mathcal{L}, \mathcal{U}]$ for $n \geq 2$, and the function*

$$f(x, y) = \frac{r + px + y}{qx + y}, \quad (x, y) \in (0, \infty)^2$$

satisfies

$$f([\mathcal{L}, \mathcal{U}] \times [\mathcal{L}, \mathcal{U}]) \subset [\mathcal{L}, \mathcal{U}]$$

Proof. Set

$$\mathcal{L} := \min \left\{ \frac{p}{q}, 1 \right\}, \quad \text{and} \quad \mathcal{U} := \max \left\{ \frac{p}{q}, 1, \frac{r + (p+1)\mathcal{L}}{(q+1)\mathcal{L}} \right\}$$

Since

$$0 \leq r + (p - q\mathcal{L})x + (1 - \mathcal{L})y \quad \text{for } (x, y) \in (\mathcal{L}, \infty)^2,$$

then $\mathcal{L}qx + \mathcal{L}y \leq r + px + qy$ for $x, y \geq \mathcal{L}$, i.e.,

$$\mathcal{L} \leq \frac{r + px + y}{qx + y}, \quad (x, y) \in (0, \infty)^2$$

From the definition of $\mathcal{U}$ we have

$$-r + (q\mathcal{U} - p)\mathcal{L} + (\mathcal{U} - 1)\mathcal{L} \geq 0$$

Write $x, y \in [\mathcal{L}, \infty)$ as $x = \mathcal{L} + v, y = \mathcal{L} + w$ for $v, w \in [0, \infty)$. Then for $v, w \in [0, \infty)$,

$$\begin{aligned}
& -r + (q\mathcal{U} - p)x + (\mathcal{U} - 1)y \\
& = -r + (q\mathcal{U} - p)(\mathcal{L} + v) + (\mathcal{U} - 1)(\mathcal{L} + w) \\
& = -r + (q\mathcal{U} - p)\mathcal{L} + (\mathcal{U} - 1)\mathcal{L} + (q\mathcal{U} - p)v + (\mathcal{U} - 1)w \\
& \geq 0
\end{aligned}$$

that is,

$$\frac{r + px + y}{qx + y} \leq \mathcal{U} \quad \text{for } x, y \in [\mathcal{L}, \infty)$$

□

Inspection of the proof of Proposition 1 reveals that, given that we have Lemma 17, the conclusion of the proposition is true concerning Eq.(3-2). The statement is given next.

Proposition 3 *At least one of the following statements is true:*

- (A) *Every solution to Eq.(3-2) converges to the equilibrium.*
- (B) *There exist m^*, M^* with $L \leq m^* < M^*$ s.t.*
 - (i) *$[m^*, M^*]$ is an invariant interval for Eq.(3-2), i.e., $f([m^*, M^*] \times [m^*, M^*]) \subset [m^*, M^*]$.*
 - (ii) *Every solution to Eq.(3-2) eventually enters $[m^*, M^*]$.*
 - (iii) *$f(x, y)$ is coordinate-wise strictly monotonic on $[m^*, M^*]^2$.*

The proof of Theorem 3 may be reproduced here in its entirety with the only change being the elimination of the case $r < 0$, which presently does not apply. Everything else in the proof applies to Eq.(3-2). The proof of Theorem 4 is complete.

7 Appendix: Computer Algebra System Code

```

g[{x_, y_}] = 
$$\frac{(1+x)(1+y)(u^2 - u + x + y)}{xy};$$


f[{x_, y_}] = 
$$\frac{(b+1)u^2 - (g+1)u + x + gy}{bx + y};$$


T[{x_, y_}] = {f[{x, y}], x};

DELTA2 = g[{x, y}] - g[T[T[{x, y}]]];

ruleb = b → 
$$\frac{z}{z+1};$$


ruleg = g → Factor[
$$\frac{\frac{1}{b}s + b}{s+1}$$
 /. ruleb];

ruleu = u → Factor[
$$\frac{g+1}{b+1} + t$$
 /. ruleg /. ruleb];

numD2 = Numerator[Together[DELTA2]];

num1D2vk = Numerator[Together[numD2 /. {x → u + v + k, y → u + v}]];

list1D2vk = Numerator[Together[num1D2vk /. {ruleu, ruleg, ruleb}]] /. Plus → List;

min21 = Min[list1D2vk /. {z → 1, s → 1, t → 1, v → 1, k → 1}];

num2D2wk = Numerator[Together[numD2 /. {x → u + v, y → u + v + k}]];

list2D2wk = Numerator[Together[num2D2wk /. {ruleu, ruleg, ruleb}]] /. Plus → List;

min22 = Min[list2D2wk /. {z → 1, s → 1, t → 1, v → 1, k → 1}];

num22 = Numerator[Together[numD2 /. {x → u / (w + 1), y → u / (v + 1)}]];

num2D2vk = Expand[num22 /. {v → w + k}];

list2D2vk = Numerator[Together[num2D2vk /. {ruleu, ruleg, ruleb}]] /. Plus → List;

min23 = Min[list2D2vk /. {z → 1, s → 1, t → 1, w → 1, k → 1}];

num2D2wk = Expand[num22 /. {w → v + k}];

list2D2wk = Numerator[Together[num2D2wk /. {ruleu, ruleg, ruleb}]] /. Plus → List;

min24 = Min[list2D2wk /. {z → 1, s → 1, t → 1, v → 1, k → 1}];

Print["Minimal Coefficients:", {min21, min22, min23, min24}];

```

Table 1: Mathematica code needed to do the calculations in Claim 3. Here we define the functions g , f and T , as well as the expression DELTA2. The reparametrizations indicated in the proof of Claim 3 for the case $g > b$ are defined as substitution rules. To verify the positive sign of a polynomial of nonnegative variables $z, s, \dots$, we form a list with the terms of the polynomial, and then substitute the number 1 for the variables in order to extract the smallest coefficient. This input was tested on Mathematica Version 5.0 [19].

```

Tz[{x_, y_}] = Together[T[{x, y}] /. {B →  $\frac{1}{1+s+t}$ , g →  $\frac{1}{1+s+t+r}$ , u → 1+s}]];

DELTA3 = Together[g[{x, y}] - g[Tz[Tz[Tz[{x, y}]]]]];

numDELTA3 = Numerator[step3];

numDELTA32 = Numerator[Together[numDELTA3 /. {x → 1+s+v, y → 1+s+w}]];

numDELTA33 = Expand[numDELTA32];

numDELTA33vk = Expand[numDELTA33 /. w → v+k];

min31 = Min[numDELTA33vk /. Plus → List /. {s → 1, t → 1, r → 1, v → 1, k → 1}];

numDELTA33wk = Expand[numDELTA33 /. v → w+k];

min32 = Min[numDELTA33wk /. Plus → List /. {s → 1, t → 1, r → 1, w → 1, k → 1}];

numDELTA32 = Numerator[Together[numDELTA3 /. {x →  $\frac{1+s}{1+v}$ , y →  $\frac{1+s}{1+w}$ }]];

numDELTA33 = Expand[numDELTA32];

numDELTA33vk = Expand[numDELTA33 /. w → v+k];

min33 = Min[numDELTA33vk /. Plus → List /. {s → 1, t → 1, r → 1, v → 1, k → 1}];

numDELTA33wk = Expand[numDELTA33 /. v → w+k];

min34 = Min[numDELTA33wk /. Plus → List /. {s → 1, t → 1, r → 1, w → 1, k → 1}];

Print["Minimal Coefficients:", {min31, min32, min33, min34}];

```

Table 2: Mathematica code needed to do the calculations in Claim 3 when $g \leq b$. The functions g , f and T are defined as before (not shown). This input was tested on Mathematica Version 5.0 [19].

References

- [1] A. M. Amleh, E. Camouzis, G. Ladas, On Second-Order Rational Difference Equations, Part 1, Journal of Difference Equations and Applications, 13(2007), 969-1004.
- [2] A. M. Amleh, E. Camouzis, G. Ladas, On Second-Order Rational Difference Equations, Part 2, Journal of Difference Equations and Applications, 14(2008), 215-228.
- [3] S. Basu and O. Merino, Global Dynamics of Second Order Rational Equations When Prime Period-Two Solutions Exist.
- [4] A. Brett and M. Kulenović, Global Asymptotic Behavior of $x_{n+1} = (px_n + x_{n-1})/(r + qx_n + x_{n-1})$, Advances in Difference Equations, 4(2007), Article ID 41541 pp. 22.
- [5] E. Camouzis and G. Ladas, *Dynamics of Third Order Rational Difference Equations*, Chapman and Hall/CRC press, 2008.
- [6] E. A. Grove, G. Ladas. Periodicities in Nonlinear Difference Equations. Chapman and Hall/CRC, Boca Raton, 2005.
- [7] R. B. Holmes, *Geometric Functional Analysis and its Applications*, Springer-Verlag New York Inc., 1975.
- [8] V. L. Kocic and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993.

- [9] M. R. S. Kulenović, Invariants and related Liapunov functions for difference equations, *Appl. Math. Lett.* 13(2000), 1–8.
- [10] M. R. S. Kulenović, personal communication.
- [11] M. R. S. Kulenović and G. Ladas, *Dynamics of Second Order Rational Difference Equations*, Chapman & Hall/CRC, Boca Raton, London, 2001.
- [12] M. R. S. Kulenović, G. Ladas, L. F. Martins, and I. W. Rodrigues, The Dynamics of $x_{n+1} = (\alpha + \beta x_n)/(A + B x_n + C x_{n-1})$ Facts and Conjectures, *Computers and Mathematics with Applications* 45(2003), 1087–1099.
- [13] M. R. S. Kulenović, G. Ladas, and W. S. Sizer, On the recursive sequence $x_{n+1} = \frac{\alpha x_n + \beta x_{n-1}}{\gamma x_n + C x_{n-1}}$, *Math. Sci. Res. Hot-Line* 2(1998), 1–16.
- [14] M. R. S. Kulenović and O. Merino, A Global Attractivity Result for Maps with Invariant Boxes, *Discrete Cont. Dynamical Syst. Ser. B* 6 (2006), 97–110.
- [15] M. R. S. Kulenović and O. Merino, *Discrete Dynamical Systems and Difference Equations with Mathematics*, Chapman and Hall/CRC Press, Boca Raton, 2002.
- [16] M. R. S. Kulenović and O. Merino, *Discrete Dynamical Systems and Difference Equations with Mathematics*, Chapman & Hall/CRC Press, Boca Raton, 2002.
- [17] G. Ladas, Invariants for generalized Lyness equations, *J. Differ. Equations Appl.* 1(1995), 209–214.
- [18] G. Ladas, On the recursive sequence $x_{n+1} = (\alpha + \beta x_n + \gamma x_{n-1})/(A + B x_n + C x_{n-1})$, *J. Differ. Equations Appl.* 1(1995), 317–321.
- [19] Wolfram Research, Inc., Mathematica, Version 5.0, Champaign, IL (2005).
- [20] L. Hu, W. Li, S. Stević, Global asymptotic stability of a second order rational difference equation. *J. Difference Equ. Appl.* 14 (2008), no. 8, 779–797. 39A11
- [21] O. Merino. Global Attractivity of the Equilibrium of a Difference Equation: An Elementary Proof Assisted by Computer Algebra System. Submitted for publication. November 2008. 10 pp.
- [22] E. C. Zeeman, Geometric Unfolding of a Difference Equation. 1996. Unpublished.